

Clinical Policy: Methoxy polyethylene glycol-epoetin beta (Mircera)

Reference Number: IL.PHAR.238

Effective Date: 4.8.20

Last Review Date: 5.6.24

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Methoxy polyethylene glycol-epoetin beta (Mircera[®]) is an erythropoiesis-stimulating agent (ESA).

FDA Approved Indication(s)

Mircera is indicated for the treatment of anemia associated with chronic kidney disease (CKD) in:

- Adult patients on dialysis and patients not on dialysis
- Pediatric patients 5 to 17 years of age on hemodialysis who are converting from another ESA after their hemoglobin level was stabilized with an ESA

Limitation(s) of use:

- Mircera is not indicated and is not recommended for use:
 - In the treatment of anemia due to cancer chemotherapy
 - As a substitute for red blood cell (RBC) transfusions in patients who require immediate correction of anemia.
- Mircera has not been shown to improve symptoms, physical functioning or health-related quality of life.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Mircera is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Anemia of Chronic Kidney Disease (must meet all):

1. Diagnosis of anemia of CKD, and member meets one of the following (a or b):
 - a. Age \geq 18 years (dialysis status is irrelevant);
 - b. Age \geq 5 years to \leq 17 years, on hemodialysis, and will be converting from another ESA agent (e.g., epoetin alfa, darbepoetin alfa)
2. Prescribed by or in consultation with a hematologist or nephrologist;
3. Adequate iron stores as indicated by current (within the last 3 months) serum ferritin level \geq 100 mcg/L or serum transferrin saturation \geq 20%;
4. Pretreatment hemoglobin $<$ 10 g/dL;

5. Failure of Epogen or Procrit unless contraindicated or clinically significant adverse effects are experienced;
6. Dosing interval does not exceed one of the following (a or b):
 - a. Adults: SC or IV once every two weeks;
 - b. Pediatrics: IV once every four weeks.

Approval duration:

Medicaid – 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Anemia of Chronic Kidney Disease (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. Failure of Epogen or Procrit unless contraindicated or clinically significant adverse effects are experienced;
4. Current hemoglobin ≤ 12 g/dL;
5. Adequate iron stores as indicated by current (within the last 3 months) serum ferritin level ≥ 100 mcg/L or serum transferrin saturation $\geq 20\%$;
6. Dosing interval does not exceed one of the following (a or b):
 - a. Adults: SC or IV once every two weeks;
 - b. Pediatrics: IV once every four weeks.

Approval duration:

Medicaid – 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):

Indication	Dosing Regimen	Maximum Dose
	<p>Conversion from another ESA: dosed SC or IV once monthly or once every two weeks based on total weekly epoetin alfa or darbepoetin alfa dose at time of conversion</p> <p>Pediatric patients with CKD on hemodialysis Conversion from another ESA: dosed IV once every four weeks based on total weekly epoetin alfa or darbepoetin alfa dose at time of conversion.</p>	

VI. Product Availability

Injection (single-dose prefilled syringe): 30, 50, 75, 100, 120, 150, 200, or 250 mcg in 0.3 mL solution; 360 mcg in 0.6 mL solution

VII. References

1. Mircera Prescribing Information. South San Francisco, CA: Genentech USA; June 2018. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/125164s0871blcorrection.pdf. Accessed January 9, 2024.
2. Kidney Disease Improving Global Outcomes (KDIGO) Clinical Practice Guideline for Anemia in Chronic Kidney Disease. Official Journal of the International Society of Nephrology – Kidney International Supplements August 2012. 2(4): 279-335.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J0887	Injection, epoetin beta, 1 microgram, (for ESRD on dialysis)
J0888	Injection, epoetin beta, 1 microgram, (for Non ESRD use)

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created, adapted CP.PHAR.238 Methoxy polyethylene glycol-epoetin beta (Mircera) for migration to HFS PDL.	4.8.20	
2Q 2021 for request for continuation add Failure of Epogen or Procrit unless contraindicated or clinically significant adverse effects are experienced; references are updated and reviewed	5.24.21	
2Q 2022 annual review: references reviewed and updated	5.23.22	

Reviews, Revisions, and Approvals	Date	P&T Approval Date
2Q 2023 annual review: no significant changes; template changes applied to other diagnoses/indications and continued therapy section; references reviewed and updated.	4.18.23	
2Q 2024 annual review: added requirement for continuation requests that current hemoglobin \leq 12 g/dL; updated dosing in Section V to include 1.2 mcg/kg once monthly dosing option for adult patients with CKD not on dialysis; references reviewed and updated.	5.6.24	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to

recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2020 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.